



TOTALLY THERE FOR YOU



PPN / SELECT - NON-GRANDFATHERED

CALENDAR YEAR DEDUCTIBLE					
Per Individual Contract	\$300	Per Individual Contract	\$600	Per Individual Contract	\$900
Per Family Contract	\$600	Per Family Contract	\$1,200	Per Family Contract	\$1,800
OUT-OF-POCKET MAXIMUM (INCLUDES DEDUCTIBLES AND COINSURANCE)					
Per Individual Contract	\$300	Per Individual Contract	\$2,000	Per Individual Contract	\$7,500
Per Family Contract	\$600	Per Family Contract	\$4,000	Per Family Contract	\$15,000
PERCENTAGE COINSURANCE PAID BY MEMBER					
	0%		20% of Cofinity (PPOM) Rate		40% of Charges

PHYSICIAN SERVICES / PREVENTIVE SERVICES

Primary care office visits	\$2 Co-pay	\$25 Co-pay	\$40 Co-pay	Deductible and Co-pay
Specialist office visits	\$2 Co-pay	\$25 Co-pay	\$40 Co-pay	Deductible and Co-pay
Allergy injections	\$2 Co-pay	\$25 Co-pay	\$40 Co-pay	Deductible and Co-pay
Chiropractic care (20 visits per year)	\$2 Co-pay	\$25 Co-pay	\$40 Co-pay	Deductible and Co-pay
Hearing and vision screening	Covered	Covered	\$40 Co-pay	Deductible and Co-pay
Immunizations (pediatric)	Covered	Covered	\$40 Co-pay	Deductible and Co-pay
Well child care	Covered	Covered	\$40 Co-pay	Deductible and Co-pay
Annual well woman visit	Covered	Covered	\$40 Co-pay	Deductible and Co-pay
Nutritional counseling and education	Covered	Covered	\$40 Co-pay	Deductible and Co-pay
Health education and counseling	Covered	Covered	\$40 Co-pay	Deductible and Co-pay
US Preventive Services Task Force A&B Recommendations	Covered (Back Page)	Covered (Back Page)	\$40 Co-pay	Deductible and Co-pay

MATERNITY SERVICES

Prenatal & postnatal care	\$2 Co-pay (one time Co-pay)	\$25 Co-pay (one time Co-pay)	\$40 Co-pay (one time Co-pay)	Deductible and Coinsurance
Delivery in hospital	Subject to Deductible		Deductible and Coinsurance	Deductible and Coinsurance
Well baby care in hospital	Covered		Coinsurance	Coinsurance

INPATIENT HOSPITAL SERVICES

Unlimited days in a semi-private room; surgery, all physicians and other ancillary services; related drug therapy; lab tests and x-rays	Subject to Deductible		Deductible and Coinsurance	Deductible and Coinsurance
---	-----------------------	--	----------------------------	----------------------------

OUTPATIENT PROCEDURES

Surgery and all invasive procedures conducted in any outpatient setting, including physicians and other ancillary services; related drug therapy; lab tests and x-rays	Subject to Deductible		Deductible and Coinsurance	Deductible and Coinsurance
--	-----------------------	--	----------------------------	----------------------------

EMERGENCY MEDICAL SERVICES

Physician and hospital emergency room services (Co-pay waived if admitted)	\$125 Co-pay	\$125 Co-pay	\$125 Co-pay	\$125 Co-pay
Ambulance services (when medically necessary)	\$75 Co-pay	\$75 Co-pay	\$75 Co-pay	\$75 Co-pay

AFTER HOURS MEDICAL SERVICES

Participating after-hours care centers (Urgent Care)	\$25 Co-pay	\$25 Co-pay	\$50 Co-pay	Deductible and Co-pay
--	-------------	-------------	-------------	-----------------------

DIAGNOSTIC & THERAPEUTIC SERVICES

Outpatient lab tests; diagnostic x-rays; and radiation therapy at a hospital	Deductible		Deductible and Coinsurance	Deductible and Coinsurance
Chemotherapy	Deductible		Deductible and Coinsurance	Deductible and Coinsurance
Physical, occupational and speech therapy (limited to 45 days per year)	\$2 Co-pay	\$25 Co-pay	\$40 Co-pay	Deductible and Co-pay
Non-preventative diagnostics including lab, x-ray, pathology, and special diagnostics in a non-hospital setting	Deductible		Deductible and Coinsurance	Deductible and Coinsurance
Mammograms	Covered		\$40 Co-pay	Deductible and Co-pay

BEHAVIORAL HEALTH CARE

Outpatient treatment	Covered		\$40 Co-pay	Not Covered
Inpatient psychiatric hospital services	Deductible		Deductible and Coinsurance	Not Covered

SUBSTANCE ABUSE TREATMENT

Outpatient Care	Covered		\$40 Co-pay	Not Covered
Intermediate Care	Deductible		Deductible and Coinsurance	Not Covered

OTHER SERVICES

Home Health Care (limited to 100 visits/year)	Covered		Not Covered	Not Covered
Hospice Care	Covered		Subject to Deductible and Coinsurance	Deductible and Coinsurance

SKILLED NURSING CARE FACILITY

limited to 120 days per year	Subject to Total Health Care USA Rider			
	Covered		Not Covered	Not Covered

DURABLE MEDICAL EQUIPMENT / PROSTHETIC DEVICES

Covered when medically necessary	Subject to Total Health Care USA Rider			
	Covered		Not Covered	Not Covered

HEARING SERVICES

Subject to Total Health Care USA Rider				
Hearing exam and hearing aid testing	Covered		Not Covered	Not Covered
Hearing aid (limited to one every three years)	Covered		Not Covered	Not Covered

VISION SERVICES

Subject to Total Health Care USA Rider				
Eye exam (limited to one year)	Covered		Not Covered	Not Covered
Eyeglasses (limited to one pair every two years)	Covered		Not Covered	Not Covered

PRESCRIPTION DRUG SERVICES

Subject to Total Health Care USA Rider				
Formulary medications prescribed by a THC participating provider and through a THC participating pharmacy	\$10/Generic \$20/Brand Name*			
*When NO Generic equivalent is available				



US Preventive Services Task Force

A & B Recommendations

Abdominal aortic aneurysm screening: men

The USPSTF recommends one-time **screening** for abdominal aortic aneurysm by ultrasonography in men aged 65 to 75 who have ever smoked.

Alcohol misuse counseling

The USPSTF recommends **screening and behavioral counseling interventions** to reduce alcohol misuse by adults, including pregnant women, in primary care settings.

Anemia screening: pregnant women

The USPSTF recommends routine **screening** for iron deficiency anemia in asymptomatic pregnant women.

Aspirin to prevent CVD: men

The USPSTF **recommends the use of aspirin** for men age 45 to 79 years when the potential benefit due to a reduction in myocardial infarctions outweighs the potential harm due to an increase in gastrointestinal hemorrhage.

Aspirin to prevent CVD: women

The USPSTF **recommends the use of aspirin** for women age 55 to 79 years when the potential benefit of a reduction in ischemic strokes outweighs the potential harm of an increase in gastrointestinal hemorrhage.

Bacteriuria screening: pregnant women

The USPSTF recommends **screening** for asymptomatic bacteriuria with urine culture for pregnant women at 12 to 16 weeks gestation or at the first prenatal visit, if later.

Blood pressure screening

The USPSTF recommends **screening** for high blood pressure in adults aged 18 and older.

BRCA screening, counseling

The USPSTF recommends that women whose family history is associated with an increased risk for deleterious mutations in BRCA1 or BRCA2 genes be **referred for genetic counseling and evaluation** for BRCA testing.

Breast cancer preventive medication

The USPSTF recommends that clinicians **discuss** chemoprevention with women at high risk for breast cancer and at low risk for adverse effects of chemoprevention. Clinicians should inform patients of the potential benefits and harms of chemoprevention.

Breast cancer screening

The USPSTF recommends **screening** mammography for women, with or without clinical breast examination, every 1-2 years for women aged 40 and older.

Breastfeeding counseling

The USPSTF recommends **interventions** during pregnancy and after birth to promote and support breastfeeding.

Cervical cancer screening

The USPSTF strongly recommends **screening** for cervical cancer in women who have been sexually active and have a cervix.

Chlamydia infection screening: non-pregnant women

The USPSTF recommends **screening** for **chlamydia** infection for all sexually active non-pregnant young women aged 24 and younger and for older non-pregnant women who are at increased risk.

Chlamydia infection screening: pregnant women

The USPSTF recommends **screening** for **chlamydia** infection for all pregnant women aged 24 and younger and for older pregnant women who are at increased risk.

Cholesterol abnormalities screening: men 35 and older

The USPSTF strongly recommends **screening** men aged 35 and older for lipid disorders.

Cholesterol abnormalities screening: men younger than 35

The USPSTF recommends **screening** men aged 20 to 35 for lipid disorders if they are at increased risk for coronary heart disease.

Cholesterol abnormalities screening: women 45 and older

The USPSTF strongly recommends **screening** women aged 45 and older for lipid disorders if they are at increased risk for coronary heart disease.

Cholesterol abnormalities screening: women younger than 45

The USPSTF recommends **screening** women aged 20 to 45 for lipid disorders if they are at increased risk for coronary heart disease.

Colorectal cancer screening

The USPSTF recommends **screening** for colorectal cancer using fecal occult blood testing, sigmoidoscopy, or colonoscopy, in adults, beginning at age 50 years and continuing until age 75 years. The risks and benefits of these screening methods vary.

Dental cares chemoprevention: preschool children

The USPSTF recommends that primary care clinicians **prescribe** oral fluoride supplementation at currently recommended doses to preschool children older than 6 months of age whose primary water source is deficient in fluoride.

Depression screening: adolescents

The USPSTF recommends **screening** of adolescents (12-18 years of age) for major depressive disorder when systems are in place to assure accurate diagnosis, psychotherapy (cognitive-behavioral or interpersonal), and follow-up.

Depression screening: adults

The USPSTF recommends **screening** adults for depression when staff-assisted depression care supports are in place to assure accurate diagnosis, effective treatment, and follow-up.

Diabetes screening

The USPSTF recommends **screening** for type 2 diabetes in asymptomatic adults with sustained blood pressure (either treated or untreated) greater than 135/80 mm Hg.

Folic acid supplementation

The USPSTF recommends that all women planning or capable of pregnancy take a daily supplement containing 0.4 to 0.8 mg (400 to 800 µg) of folic acid.

Gonorrhea prophylactic medication: newborns

The USPSTF strongly recommends prophylactic ocular topical medication for all newborns against gonococcal ophthalmia neonatorum.

Gonorrhea screening: women

The USPSTF recommends that clinicians **screen** all sexually active women, including those who are pregnant, for gonorrhea infection if they are at increased risk for infection (that is, if they are young or have other individual or population risk factors).

Healthy diet counseling

The USPSTF recommends intensive behavioral dietary **counseling** for adult patients with hyperlipidemia and other known risk factors for cardiovascular and diet-related chronic disease. Intensive counseling can be delivered by primary care clinicians or by referral to other specialists, such as nutritionists or dietitians.

Hearing loss screening: newborns

The USPSTF recommends **screening** for hearing loss in all newborn infants.

Hemoglobinopathies screening: newborns

The USPSTF recommends **screening** for sickle cell disease in newborns.

Hepatitis B screening: pregnant women

The USPSTF strongly recommends **screening** for hepatitis B virus infection in pregnant women at their first prenatal visit.

HIV screening

The USPSTF strongly recommends that clinicians **screen** for human immunodeficiency virus (HIV) all adolescents and adults at increased risk for HIV infection.

Hypothyroidism screening: newborns

The USPSTF recommends **screening** for congenital hypothyroidism in newborns.

Iron supplementation in children

The USPSTF recommends routine iron supplementation for asymptomatic children aged 6 to 12 months who are at increased risk for iron deficiency anemia.

Obesity screening and counseling: adults

The USPSTF recommends that clinicians **screen** all adult patients for obesity and offer **intensive counseling and behavioral interventions** to promote sustained weight loss for obese adults.

Obesity screening and counseling: children

The USPSTF recommends that clinicians **screen** children aged 6 years and older for obesity and **offer them or refer them to comprehensive, intensive behavioral interventions** to promote improvement in weight status.

Osteoporosis screening: women

The USPSTF recommends that women aged 65 and older be **screened** routinely for osteoporosis. The USPSTF recommends that **routine screening** begin at age 60 for women at increased risk for osteoporotic fractures.

PKU screening: newborns

The USPSTF recommends **screening** for phenylketonuria (PKU) in newborns.

Rh incompatibility screening: first pregnancy visit

The USPSTF strongly recommends Rh (D) blood typing and antibody testing for all pregnant women during their first visit for pregnancy-related care.

Rh incompatibility screening: 24-28 weeks gestation

The USPSTF recommends repeated Rh (D) antibody **testing** for all unsensitized Rh (D)-negative women at 24-28 weeks gestation, unless the biological father is known to be Rh (D)-negative.

STIs counseling

The USPSTF recommends high-intensity **behavioral counseling** to prevent sexually transmitted infections (STIs) for all sexually active adolescents and for adults at increased risk for STIs.

Tobacco use counseling: non-pregnant adults

The USPSTF recommends that clinicians **ask** all adults about tobacco use and provide tobacco cessation **interventions** for those who use tobacco products.

Tobacco use counseling: pregnant women

The USPSTF recommends that clinicians ask all pregnant women about tobacco use and provide augmented, pregnancy-tailored **counseling** to those who smoke.

Syphilis screening: non-pregnant persons

The USPSTF strongly recommends that clinicians **screen** persons at increased risk for syphilis infection.

Syphilis screening: pregnant women

The USPSTF recommends that clinicians **screen** all pregnant women for syphilis infection.

Visual acuity screening in children

The USPSTF recommends **screening** to detect amblyopia, strabismus, and defects in visual acuity in children younger than age 5 years.